

IOWA STATE UNIVERSITY

Digital Repository

Integrated Crop Management News

Extension and Experiment Station Publications

5-8-2018

Soybean Aphid Egg Hatch Predicted in Northern Iowa

Erin Hodgson

Iowa State University, ewh@iastate.edu

Follow this and additional works at: <https://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), and the [Agriculture Commons](#)

Recommended Citation

Hodgson, Erin, "Soybean Aphid Egg Hatch Predicted in Northern Iowa" (2018). *Integrated Crop Management News*. 2498.
<https://lib.dr.iastate.edu/cropnews/2498>

The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.

Soybean Aphid Egg Hatch Predicted in Northern Iowa

Abstract

Iowa's most significant soybean insect pest, soybean aphid, has host-alternating biology. This species has multiple, overlapping generations on soybean in the summer and moves to buckthorn in the winter. Fall migration to buckthorn is based on senescing soybean, and decreasing temperatures and photoperiod. For the majority of the year, soybean aphids are cold-hardy eggs near buckthorn buds (Photo 1). As spring temperatures warm up, soybean aphid eggs hatch and produce a few generations on buckthorn before moving to soybean (Photo 2). Tilmon et al. (2011) goes into more detail about the life cycle and biology of soybean aphid.

Disciplines

Agricultural Science | Agriculture

IOWA STATE UNIVERSITY

Extension and Outreach

Integrated Crop Management

Soybean Aphid Egg Hatch Predicted in Northern Iowa

May 8, 2018

Iowa's most significant soybean insect pest, soybean aphid, has host-alternating biology. This species has multiple, overlapping generations on soybean in the summer and moves to buckthorn in the winter. Fall migration to buckthorn is based on senescing soybean, and decreasing temperatures and photoperiod. For the majority of the year, soybean aphids are cold-hardy eggs near buckthorn buds (Photo 1). As spring temperatures warm up, soybean aphid eggs hatch and produce a few generations on buckthorn before moving to soybean (Photo 2). Tilmon et al. (2011) goes into more detail about the life cycle and biology of soybean aphid.



Photo 1. Sexual females deposit eggs near buckthorn buds in the fall. Photo by David Voegtlin.



Photo 2. There are a few wingless generations produced on buckthorn before the spring migration to soybean every year. Photo by Chris DiFonzo (*Bugwood.org*).

For many aphids that overwinter as an egg, hatching often happens when the host resumes spring growth. This makes biological sense because the aphids feed on phloem from actively-growing tissue. If egg hatch happens too soon, they can suffer mortality from starvation. Research has confirmed soybean aphid egg hatch happens around buckthorn bud swell. Bahlai et al. (2007) developed a model to predict soybean aphid egg hatch based on accumulating degree days. They adjusted the model to include ambient air temperatures and solar radiation. Soybean aphid egg hatch occurs between 147-154 degree days (base 50°F) and buckthorn bud swell happens shortly after that (165-171 degree days). Based on air temperatures in 2018, we expect egg hatch is occurring in northern Iowa, where most of the buckthorn in Iowa is located (Figure 1).




Figure 1. Accumulated growing degree

days (base 50°F) in Iowa from January 1 – May 7, 2018. Map courtesy of Iowa Environmental Mesonet, ISU Department of Agronomy.

References

Bahlai, C. A., J. A. Welsman, A. W. Schaafsma, and M. K. Sears. 2007. Development of soybean aphid on its primary overwintering host. *Environmental Entomology* 36: 998-1006.

Tilmon, K. J., E. W. Hodgson, M. E. O'Neal, and D. W. Ragsdale. 2011. Biology of the soybean aphid in the United States. *Journal of Integrated Pest Management* 2: A1-A7.

Links to this article are strongly encouraged, and this article may be republished without further permission if published as written and if credit is given to the author, Integrated Crop Management News, and Iowa State University Extension and Outreach. If this article is to be used in any other manner, permission from the author is required. This article was originally published on May 8, 2018. The information contained within may not be the most current and accurate depending on when it is accessed.

Category: [Crop Production](#) [Insects and Mites](#)

Crop:

[Soybean](#)

Tags: [Soybean](#) [pest](#) [IPM](#) [scouting](#)

Author:



Erin Hodgson Associate Professor

Dr. Erin Hodgson started working in the Department of Entomology at Iowa State University in 2009. She is an associate professor with extension and research responsibilities in corn and soybeans. She has a general background in integrated pest management (IPM) for field crops. Dr. Hodgson's curre...